

Gold Coast Primary Health Network Needs Assessment 2022



Crisis support and
suicide prevention

phn
GOLD COAST

An Australian Government Initiative

Crisis support and suicide prevention

Local health needs and service issues

- Suicidal people often visit primary care providers in the weeks or days before suicide but often their suicide risk is not identified.
- Limited supports are available for people in distress who end up in Emergency Department by default or on a mental health trajectory.
- Carers are at the frontline of suicide prevention and many sit daily with this risk and responsibility but may not have any training or skills to equip them for this.
- When challenges occur during a crisis, it is often at the points of intersection between different sectors.
- Service providers do not always know what the best evidence-based treatments are for people experiencing suicidal thoughts and behaviours or how to access local services and supports.
- Organisations can be very risk averse due to a culture of blame. People in suicidal crisis can be passed back and forth between organisations without receiving the care they need.
- Many mental health clinicians do not have specific training in suicide prevention or know what the best evidence-based treatments are for people experiencing suicidal thoughts and behaviours.
- Many people in the community lack the confidence and skills to address people in suicidal distress or crisis.
- People with a lived experience of suicide have the potential, to inform, influence and enhance local suicide prevention solutions but may lack the confidence, skills and readiness to participate.
- Confidentiality and legal concerns along with lack of formal arrangements limit information sharing between providers.

Key findings

- Gold Coast Primary Health Network (GCPHN) suicide rate is consistent with the state rate, while greater than the national rate.
- The Gold Coast has one of the busiest Emergency Departments in Queensland, and a large percentage of these are people presenting with mental health issues.
- Males accounted for 65% of suspected deaths by suicide on the Gold Coast in 2017-2019, compared to 35% for females in the same period.
- Mental illness, including depression, and trauma are associated with a large portion of suicide attempts.
- Suicide prevention is the most common cause for referral by General Practitioners (GP) to the Psychological Services Program.
- National data indicates the lesbian, gay, bisexual, transgender, intersex, queer, asexual, pansexual and others communities, Aboriginal and Torres Strait Islanders, and culturally and linguistically diverse community is particularly vulnerable.
- Services that support people struggling with relationship and family breakdowns, financial problems and bereavement are essential elements of the suicide prevention system.

- A mental health crisis is not synonymous with mental illness. A prolonged mental health crisis can however lead to a mental illness.

Prevalence, service usage and other data

Deaths by suicide

Suicide was the leading cause of death for young Queenslanders in 2020 with 126 deaths among people aged 15-24 years¹. It was also the leading cause of death for people aged 25-34 years with 139 deaths and 35-44-year-olds with 149 deaths in Queensland.

In the 2020 calendar year, there were 759 suspected suicides of Queensland residents. This is just under 15 suspected suicides for every 100,000 people. The number of male suspected suicides increased by 28 from 570 in 2019 to 598 in 2020. Female suspected suicides decreased by 26, from 187 in 2019 to 161 in 2020. Strangulation or suffocation and poisoning were the most common methods of suicide in Queensland.

Suspected suicide rates of Queensland residents have decreased since 2017 with a 2.8% decrease from 2018 to 2019 in Queensland. Male suspected suicide rates in 2020 increased from 22.9 to 24.0 per 100,000 males. For females the number decreased with 7.3 per 100,000 in 2019 to 6.6 per 100,000 in 2020. Suspected suicide numbers and rates for 2020 were highest in males aged 75 to 79 and females aged 30-34. These age cohorts differ from the age cohorts with the highest rates in 2019 of males 40-49 and females aged 45-49².

During the period 2017-2019, there were 278 suicides in the Gold Coast Primary Health Network catchment area, representing an age-standardised suicide rate of 14.5 per 100,000 people².

Table 1. Suspected suicides and suicide rates per 100,000 people, Primary Health Network, Queensland, 2017–19

PHN	Suspected Suicides	Age-standardised rate per 100,000
Northern Queensland	423	20.6
Western Queensland	35	20.0
Darling Downs and West Moreton	303	18.3
Central Queensland, Wide Bay, Sunshine Coast	424	16.8
Gold Coast	278	14.5
Brisbane North	407	13.1
Brisbane South	545	12.9

Source: Leske, S., Adam, G., Schrader, I., Catakovic, A., Weir, B., & Crompton, D. (2020). *Suicide in Queensland: Annual Report 2020*. Brisbane, Australia.

The release of the 2015-2019 leading cause of death in Australia by Australian Institute of Health and Welfare indicated suicide was the 11th leading cause of death in this period with 15,743 deaths (12.7) per 100,000)


¹ Australian Bureau of Statistics, 3303.0-Cause of Death.

² s S Leske, I Schrader, G Adam, A Catakovic, B Weir and K Kőlves, *Suicide in Queensland: annual report 2021*

in Australia. Suicide was the 8th leading cause of death on the Gold Coast in the same reporting period with 447 deaths (14.5 per 100,000)³.

As can be seen in Figure 1, the Gold Coast rate of deaths by suicide from 2015 to 2019 was above the national rate while Surfers Paradise had the largest rate of deaths by suicide with 19.2. Although the age-standardised rate of deaths by suicide identified that Ormeau-Oxenford had the second lowest rate the GCPHN region, the region did have the highest total number of deaths by suicide from 2014 to 2018 (Table 2).

Figure 1. Age-standardised rate per 100,000 people of deaths by suicide, Gold Coast SA3 regions, 2015-19



Source: Deaths in Australia/Grim MORT Books, Australian Institute of Health and Welfare, 2020

Table 2 shows the number of deaths by suicide across Gold Coast SA3 regions between 2015 and 2019, identifying the northern corridor of the Gold Coast as having the highest number of suspected suicides.

³ AIHW (Australian Institute of Health and Welfare) 2021. MORT (Mortality Over Regions and Time) books: Primary Health Network (PHN), 2015–2019. Canberra: AIHW.

Table 2. Suicides in Gold Coast PHN including SA3 regions, 2015–19


	Number of deaths
Gold Coast SA4	447
Ormeau - Oxenford	88
Nerang	61
Gold Coast - North	54
Surfers Paradise	51
Southport	45
Broadbeach - Burleigh	43
Coolangatta	39
Robina	27
Mudgeeraba - Tallebudgera	27
Gold Coast Hinterland	12

Source: AIHW (Australian Institute of Health and Welfare) 2021. MORT (Mortality Over Regions and Time) books: Statistical Area Level 3 (SA3), 2015–2019. Canberra: AIHW.

Suicide rates by age and gender

In 2018, the age group of 50-54 had the highest number people dying of suicide in the Gold Coast Hospital and Health Service compared to 2013, which the age group of 40-44 had the highest number of people dying by suicide as can be seen in Figure 2.


Figure 2. Deaths by suicide, by age groups, Gold Coast and Hospital and Health Service, 2013 and 2018



Source: Leske, S., Crompton, D., & Kölves, K. (2019). *Suicide in Queensland: Annual Report 2019*. Brisbane, Queensland, Australia: Australian Institute for Suicide Research and Prevention, Griffith University

The male rate of deaths by suicide has been decreasing on the Gold Coast in recent years while the female rate has been increasing. Males accounted for 59.4% of deaths by suicide on the Gold Coast in 2018 compared to 76.1% in 2013. In 2018, the female rate of deaths by suicide was 40.6% which has increased from 23.9% in 2013.

Figure 3. Deaths by suicide, by sex, Gold Coast and Hospital and Health Service, 2013 to 2018



Source: Leske, S., Crompton, D., & Kölves, K. (2019). *Suicide in Queensland: Annual Report 2019*. Brisbane, Queensland, Australia: Australian Institute for Suicide Research and Prevention, Griffith University

Aboriginal and Torres Strait Islander peoples

The suicide rate in Queensland Aboriginal and Torres Strait Islander peoples is twice that of the non-Indigenous population, and suicide occurs at a much younger age. Intentional self-harm is the fifth highest cause of death for Indigenous people, with males representing the vast majority (83%) of suicide deaths⁴.

Of the 757 suicides reported in 2019 in Queensland, Aboriginal and Torres Strait Islander females living in Queensland accounted (11.9%) of all female suicides while males accounted for 8.3% of all male suicides⁵. The age group of 20-24 had the highest number of suspected suicides by Aboriginal and Torres Strait Islander Queenslanders.

True suicide mortality figures in Aboriginal and Torres Strait Islander populations remain poorly understood due to incomplete data collection processes and inaccurate classification systems (see Mental Health & Suicide Aboriginal & Torres Strait Islander needs assessment).

⁴ Australian Bureau of Statistics (2018). Catalogue 3303.0—Causes of Death. Canberra, Australia

⁵ Leske, S., Adam, G., Schrader, I., Catakovic, A., Weir, B., & Crompton, D. (2020). *Suicide in Queensland: Annual Report 2020*. Brisbane, Queensland, Australia: Australian Institute for Suicide Research and Prevention, Griffith

Lesbian, gay, bisexual, transgender, intersex, queer, asexual, pansexual, and other (LGBTIQAP+)

LGBTIQAP+ are far more likely to attempt suicide than heterosexual people. LGBTIQAP+ people are between 3.5 and 14-times more likely to try and die by suicide compared to heterosexual people⁶.

Of the 757 suicides reported in 2019 in Queensland, 36 (1.5% of all) suspected suicides by persons identifies as LGBTIQAP+⁵. Australian Bureau of Statistics data indicates a heightened risk of poor mental health that may lead to suicidal behaviour in LGBTIQAP+ communities⁷. This increased risk of poor mental health and suicidality among LGBTIQAP+ people are not attributable to sexuality, sex, or gender identity, but rather due to experiences of discrimination and exclusion⁸.

One in six young LGBTIQAP+ people have attempted suicide and one third have harmed themselves. 16% of LGBTIQAP+ Australians aged between 16 and 27 have attempted suicide and 33% have self-harmed⁹. Looking at transgender young people, around 3 in every 4 transgender young people have experienced anxiety or depression, 4 out of 5 transgender young people have ever engaged in self-harm and almost 1 in 2 (48%) have attempted suicide¹⁰.

Culturally and linguistically diverse (CALD) populations

Australia's CALD communities have diverse views of suicide and suicidal thinking, and vary in the way that their community, family, and friends respond to suicide. Multicultural differences, past trauma and experiences of discrimination are acknowledged and related to effective suicide prevention strategy. Limited data is available on this group although stigma around mental health and the topic of suicide, as well as language barriers and the difficulty of maintaining privacy and confidentiality can affect people in CALD communities.

Prevalence of life events

There are multiple factors recognised as contributing to suicidal behaviour or someone being at risk of suicide. These include personal hardship, difficult life events, poor physical and mental health such as depression and trauma, harmful substance use and previous self-harm or suicide attempts. It is important to understand these factors when considering suicide prevention.

Data from the Australian Institute for Suicide Research and Prevention identified the prevalence of life events among people who died by suicide (2013-2015). Relationship separation was the most frequently recorded life event (32.5%) among all ages and for both women and men. This was followed by financial problems (27.3%), recent or pending unemployment (19.5%).

⁶ Suicide Prevention Australia Position Statement, Suicide and self-harm among Gay, Lesbian, Bisexual and Transgender communities 2009

⁷ Skerrett, D., Kolves, K., De Leo, D (2015). Are LGBT Populations at a Higher Risk for Suicidal Behaviors in Australia? Research Findings and Implications. *Journal of Homosexuality*. Vol. 62. Issue 7.

⁸ Rosenstreich, G. (2013). *LGBTI People Mental Health and Suicide*. Revised 2nd Edition. National LGBTI Health Alliance: Sydney.

⁹ Robinson, KH, Bansel, P, Denson, N, Ovenden, G & Davies, C 2014, *Growing Up Queer: Issues Facing Young Australians Who Are Gender Variant and Sexuality Diverse*, Young and Well Cooperative Research Centre, Melbourne

¹⁰ Strauss, P., Cook, A., Winter, S., Watson, V., Wright Toussaint, D., Lin, A. (2017). *Trans Pathways: the mental health experiences and care pathways of trans young people*. Summary of results. Telethon Kids Institute, Perth, Australia

Table 3. Life events reportedly experienced by those dying by suicide in Gold Coast Hospital and Health Service, 2013 to 2015

Life event		2013	2014	2015
Relationship problems	Conflict	17.9%	14.1%	18.2%
	Separation	22.4%	29.4%	32.5%
Bereavement	Spouse	9.0%	3.5%	3.9%
	Family	6.0%	7.1%	5.2%
	Other	1.5%	1.2%	3.9%
	Multiple	1.5%	0.0%	0.0%
Conflict	Familial	6.0%	8.2%	10.4%
	Interpersonal	7.5%	3.5%	7.8%
Other	Pending legal matters	4.5%	8.2%	5.2%
	Financial problems	14.9%	20.0%	27.3%
	Recent/pending unemployment	9.0%	8.2%	19.5%
	Work/school problems	4.5%	7.1%	5.2%
	Child custody dispute	6.0%	4.7%	6.5%
	Childhood trauma	6.0%	0.0%	3.9%
	Sexual abuse	1.5%	0.0%	2.6%

Source: Leske, S., Crompton, D., & Kölves, K. (2019). *Suicide in Queensland: Annual Report 2019*. Brisbane, Queensland, Australia: Australian Institute for Suicide Research and Prevention, Griffith University.

ED Presentations

EDs are frequent places for people in mental health crisis to present, with 69,585 presentations in 2019/20 by Gold Coast residents. Of these presentations, 59% the emergency service episode is completed and discharged, 19% are admitted, 16% are admitted to short stay unit, 4% transferred to another hospital, 2% left at own risk after treatment commenced and 1% did not wait.

In general people presenting with mental health issues wait longer to be seen initially in EDs than other consumers with a similar severity of physical illness and of concern, they were twice as likely as other ED presentations to leave before their treatment and care was complete. Crisis responses do not respond well to the needs of individuals and emergency mental healthcare is frequently compared unfavorably to emergency physical care, raising issues of lack of equality.

Presentations to Gold Coast University Hospital and Robina Hospital Emergency Department for suicidal ideation between June 2019 to July 2020 was slightly above 2,000 presentations. Of these, 49% were males and 51% were females. 7% of presentations were by people identifying as Aboriginal and or Torres Strait Islanders. People aged between 20 to 29 years old had the largest rate of presentation of people for suicidal ideation with 29%, followed by people aged 7 to 19 (23%).

Intentional self-harm

Intentional self-harm is often defined as deliberately injuring or hurting oneself, with or without the intention of dying. Intentional self-harm comes in many forms, and affects people from different backgrounds, ages and lifestyles. The reasons for self-harm are different for each person and are often complex. Most people who self-harm does not go on to end their lives- but previous self-harm is a strong risk factor for suicide. Therefore, monitoring of intentional self-harm is key to suicide prevention.

As can be seen in Table 4, the GCPHN region was below the Queensland rate per 100,000 people for all intentional self-harm for all age cohorts except males aged 0-24 and females aged 65 years and over in 2019-20.

Table 4. Number and rate of intentional self-harm hospitalisations, Gold Coast and Queensland, 2019-20

		Gold Coast	Queensland	Gold Coast	Queensland
		Number		Rate per 100,000 people	
Male	0-24	103	946	102.9	112.1
	25-44	134	1,328	158.9	194.5
	45-64	83	737	110.3	120.4
	65+	24	203	48.3	53.4
	All ages	344		3,214	
Female	0-24	219	2,467	305.3	293.6
	25-44	173	1,723	243.9	251.6
	45-64	119	1,018	158.7	167.4
	65+	38	214	51.0	48.8
	All ages	549		5,422	

Source: Data tables: 2019–20 National Hospital Morbidity Database—Intentional self-harm hospitalisations

Psychological Services Program

The Psychological Services Program (PSP) provides short term psychological interventions for financially disadvantaged people with non-crisis, non-chronic, moderate mental health conditions or for people who have attempted, or at risk of suicide or self-harm. This program targets seven underserved and priority groups, including children, people at risk of homelessness and suicide prevention.

From the 1st of July 2021 to 30th June 2022 there were 1,056 referrals to the Adult Suicide Prevention Psychological Services Program (PSP) stream, leading to 4,909 sessions. Suicide prevention is by far the most common cause for referral by GPs and services users include a range of people in distress. Of those referred to the adult suicide prevention stream, 16% came from clients located in Coomera, Pimpama, and Upper Coomera followed by 15% from Labrador and Southport.

Table 5. Number of persons accessing Psychological Services Program on the Gold Coast, 2021-22

	Referrals	% of all referrals	Sessions	Sessions as % of referrals
Adult Suicide Prevention	1056	65.2%	4909	70.1%
Children	235	14.5%	849	12.1%
Aboriginal and Torres Strait Islander	111	6.9%	383	5.5%
Homeless	55	3.4%	197	2.8%
CALD	47	2.9%	248	3.5%
Perinatal	68	4.2%	215	3.1%
LGBTIQAP+	48	3.0%	197	2.8%
General (COVID19 Response)	0	0.0%	1	0.01%
Total	1,620		6,999	

Source: PIR-FIXUS

The Way Back Support Service

People who have attempted suicide or experienced a suicidal crisis often experience severe distress in the days and weeks immediately afterwards, and they are at high risk of attempting again within 12 months from being discharged. Beyond Blue developed The Way Back Service Support Service to support them through this critical risk period. The Way Back provides non-clinical, tailored support for up to three months following discharge from hospital after a suicide attempt.

The Gold Coast Way Back Service receives the largest number of referrals compared to other PHN regions. Between 1st July to 30th March 2021, 432 people accessed the service which exceeds the expected number of people accessing the service of 261 (166% of target).

Suicide Prevention Pathway

The Gold Coast Mental Health and Specialist Services Suicide Prevention Pathway assists patient's recovery from suicidal thoughts and behaviors. Between January and November 2019, there was a total of 1,681 placements on the Suicide Prevention Pathway (on average, 153 placements per month). This represented a total of 1,498 persons (average of 136 / month). Of these people, 84.2% were aged 18 years and over.

COVID-19 Impacts

There is much uncertainty around the medium- and long-term impacts of the COVID-19 pandemic on suicide mortality in Australia. Duration and intensity of restrictions, timeframe of economic recovering and the impact of state and federal government interventions to reduce the economic and social effects will all affect suicide mortality. As the Gold Coast is a region dependent on tourism this may have more of an impact on the Gold Coast compared to other regions depending on ongoing social distancing practices.

As previously mentioned, suicide is not influenced or caused by one factor but results from complex interaction between multiple risk factors, consequently it is difficult to understand the impact COVID-19 on suspected suicides. As can be seen in Table 6, the year-to-date comparisons for suicides from January to July in 2020 is comparable to 2019.

Table 6. Year to date comparisons for suicides from January to July, by sex, Queensland, 2015-2020

Year	Males	Females
2015	315	106
2016	290	92
2017	340	116
2018	341	89
2019	343	102
2020	352	102

Source: Leske, S., Adam, G., Schrader, I., Catakovic, A., Weir, B., & Crompton, D. (2020). *Suicide in Queensland: Annual Report 2020*. Brisbane, Queensland, Australia: Australian Institute for Suicide Research and Prevention, Griffith University

The Wesley Mission Queensland COVID-19 Recovery Service was established to provide responsive wellbeing support for people aged 16 years and over living in the GCPHN region whose wellbeing has been impacted by the ongoing effects of COVID-19. Below are some of the common presentations to the service:

- loneliness and social isolation,
- suicidal ideation,
- problems with secure housing,
- financial barrier’s such as loss of employment/struggles to secure adequate ongoing employment,
- overall anxiety and depressive presentations – low mood and lack of motivation,
- struggles with accessing services such as Centrelink and NDIS,
- loss of routine,
- grief and loss,
- difficulties in accessing appropriate services in a timely manner due to long waitlists.


In addition to the above, very early on in the rollout of this service it became apparent that schools needed support as children’s anxiety levels had increased; anecdotally home-schooling had a massive impact on some students finding it difficult to re-engage in face-to-face learning. Parents were also struggling with how to deal with the impact of COVID-19 on the mental health and wellbeing of their children.

Community approach to suicide prevention

Reducing the rate and impact of suicide in the Gold Coast community is not something any single agency or level of government can do alone. The health system plays a vital role in suicide prevention, particularly through the delivery of specialised mental healthcare. However, equally important roles are played by a wide range of social and human services, law enforcement agencies, industry bodies, education providers, private and non-government service providers, community services and workplaces. Community events can also provide people with clear opportunities to be actively involved in suicide prevention.

Recognising the need for a community approach to suicide prevention, as part of the Joint Regional Planning process a Suicide Prevention Leadership Group was formed in August 2019. This group advised on the suicide prevention components of the Joint Regional Plan and developed a more in-depth Community Action Plan for Suicide Prevention using the LifeSpan framework developed by Black Dog Institute. This framework includes nine evidence-based strategies and six overarching principles and when implemented together, this approach is predicted to reduce suicide death by 20% and suicide attempts by 30%.

Figure 4. LifeSpan: Integrated Suicide Prevention



Source: *The LifeSpan lived experience framework, Australian National University*

Through the development of the Suicide Prevention Community Action Plan, the following issues were identified, in alignment with the nine LifeSpan strategies:

1) Improving emergency and follow up care for suicidal crisis:

- A suicide attempt is the strongest risk factor for subsequent suicide.
- People who present in emergency department in crisis or for suicidal thinking or attempts often do not receive the care and support they need. This may be related to staff experience and skills to deal with suicide and crisis.
- The emergency department environment can be fast paced and traumatising environment.
- Police and ambulance may not have the level of experience/skills or time to deal with mental health related call outs effectively.
- Current resource material to support crisis and suicide is outdated to changes in the sector.

2) Using evidence-based treatment for suicidality:

- Mental illness, including depression, and trauma are associated with a large portion of suicide attempts.
- Currently it is unclear what evidence-based treatments are being delivered, by whom or what the quality of these services is.
- Mental health professionals are not aware of the latest evidence and best practice care and treatment options for suicide.
- The Gold Coast has some of the highest use of MBS billings in the country for the private sector, but it is not clear who is accessing these services, what services are available and the quality of these services.

- There is a lack of urgency for evidence-based treatment options to address suicide within the mental health sector.
- 3) Equipping primary care to identify and support people in distress:
- Primary care providers are often visited by suicidal persons in the weeks or days before suicide but due to fear, stigma or time pressures, do not receive the care they need.
 - GPs encounter numerous barriers and competing priorities which impacts GP uptake and access to suicide prevention training.
 - Traditional GP training does not necessarily equip GPs with the skills and confidence to address mental health concerns and suicidal ideation.
 - Many GPs are unaware of referral points and current best practice care and treatment.
 - Issues with GPs being able to access forms for referral pathways other than Mental healthcare Plan e.g., Psychological Service Providers.
- 4) Improving the competency and confidence of frontline workers to deal with suicidal crisis:
- Frontline workers can play a key role in de-escalating a crisis and improving safety.
 - Existing training for frontline workers (mental health services, police, paramedics, and hospital staff) may not include specific suicide prevention skills.
 - Barriers to training such as funding, time of day, endorsement and approval by workplaces can limit uptake and participation in training.
- 5) Promoting help-seeking, mental health and resilience in schools:
- Schools are overwhelmed with options and pressure from multiple bodies/sectors to include additional content in their curriculum and programs.
 - Schools are keen to support their students but often do not know how to choose quality programs or integrate programs with other student wellbeing activities and referral pathways.
 - A focus on preventing or intervening early in the progression of mental health difficulties not only benefits infants and children, but also creates a solid foundation for health outcomes later in life.
 - Training initiatives are often fragmented, parents, teachers, and young people may all receive different training, resources and information about how to respond to mental health issues and suicidal crisis resulting in fragmentation and diffusion of responsibility.
 - Currently the communication between hospitals and schools to is not being optimised to support young people post discharge and in the recovery process or to help children and youth remain engaged with school.
 - Clinicians in schools often operate in silos and at the discretion of school principals. Involvement in the planning of school activities could facilitate and enhance coordination of activities.
 - Organisations can be very risk averse due to a culture of blame. People in suicidal crisis can be passed back and forth between organisations without getting the care they need.
- 6) Training the community to respond to suicidality/Gatekeeper training:
- Many people who are experiencing suicidal thoughts communicate distress through their words or actions, but these warning signs may be missed or misinterpreted.

- Suicide prevention efforts are often fragmented and have not always been strategically planned or coordinated.
- Inconsistent approaches to increasing Mental Health and Suicide Prevention literacy across the community through workplaces.
- Many gatekeepers are in roles that might encounter people in suicidal crisis, however since this is not their primary role, they may lack skills and confidence to respond to suicidality.
- There is a lot of training available, but people are not always aware of what is available and relevant to them, this may result in duplication and inefficiency.
- There are limited evidence around which programs are most effective and relevant to local stakeholders.

7) Engaging the community and providing opportunities to be a part of change:

- Suicide prevention activity is frequently fragmented. There are opportunities to improve awareness of how we can work together better.
- Suicide prevention services and approaches need to be more culturally inclusive and responsive to diversity.
- Service eligibility criteria/thresholds often limits access to services. In addition, many services do not provide afterhours support.
- Stigma associated with suicide and help-seeking is a significant barrier to prevention. Greater acknowledgment and recognition of community suicide prevention activity is required to raise the profile of suicide prevention and postvention in a positive way.
- There is often stigma attached to mental health and suicide. Some people don't identify with these labels and will not access support for conditions that they don't relate to.
- Safe communication about suicide actively promotes help-seeking, reduces stigma and encourages collaboration.
- People do not know how to be actively involved in suicide prevention and are not always aware of opportunities or ways they can contribute.

8) Encouraging safe and purposeful media reporting:

- Representations of suicide in the media can be sensationalised/or stigmatised and unsafe leading to copycat behaviour.
- The graphic nature of news can be traumatising and cause fear and anxiety.
- People with a lived experience of suicide are often not empowered or provided with opportunities to become agents of system change or to share messages of hope and recovery with others.
- Suicide prevention activities and campaigns could be better coordinated to maximise impact.

9) Improve safety and the means of suicide:

- Currently timely (up to date) regional data is not available which limits our ability to use data to drive decision making.
- Safety plans are held by providers and individuals have to develop new safety plans with multiple providers.
- Carers are often not aware of/informed of details of safety plans and how they can support people to implement their safety plans.

Mental Health Crisis Reform

Gold Coast Health commenced the Mental Health Crisis Reform initiative in the second half of 2019 with a consideration of the Crisis Now framework¹¹, which emphasises a number of care elements including: regional or state-wide crisis call centres coordinating in real time: centrally deployed, 24/7 mobile crisis teams; short-term, “sub-acute” residential crisis stabilization programs; and essential crisis care principles and practices including recovery orientation, trauma-informed care, significant use of peer staff, a commitment to Zero Suicide/Suicide Safer Care, strong commitments to safety for consumers and staff, and collaboration with law enforcement.

Feedback was obtained through a range of stakeholder meetings held on the Gold Coast across the second half of 2019 and through 2020. There was a very positive response to the core elements as outlined in the Crisis Now framework. However, it was felt by many stakeholders that it was important to emphasise other aspects of the system and go beyond those presented within Crisis Now. There was strong feedback that any plans around crisis reform on the Gold Coast needed to have due consideration of the whole continuum of care, and to be well integrated into the community, and recovery focused. The Gold Coast Crisis Reform Framework has been developed¹² in response to the following issues identified by local stakeholders:

- There is a need for health services (physical health, mental health, alcohol and other drugs services), social services and emergency response services (e.g., police and ambulance) to work together on coordinated and strategic approaches to transforming mental health crisis care across the GCPHN region. When challenges occur during a crisis, it is often at the points of intersection of these agencies. These entities have their own points of entry, and staff with significant variation in skills, training, and experience in mental health crisis. There are complex questions regarding who takes the lead for certain situations and how does integration and communication occur.
- While principles related to best practice crisis care have been driving reform at a regional level for many years, there is a need to continue to embed these principles in our service and system, both existing and new initiatives:
 - trauma-informed care
 - lived experience and involvement of families central to all models of care
 - adopting a Journey to Zero Seclusion and Restraint
 - integrated mental health, alcohol and other drug and physical healthcare
 - culturally safe, responding to diversity
- A narrow focus on how we respond once a significant crisis has developed will not meet the needs of our community, nor will it align with a growing evidence base internationally. Only with an adequate continuum of service will we be able to prevent crises from developing or reduce likelihood of re-presentations in the future. A comprehensive system needs to include social and housing support to enable recovery and prevent a cycle of repeated crises.
- There is a need for real-time displays of data to inform rapid decision making and tracking of consumers during their crises.

¹¹ *Crisis Now: Transforming Services is Within our Reach* (National Action Alliance for Suicide Prevention: Crisis Services Task Force, 2016)

¹² Gold Coast Crisis Reform: A Strategic Approach to Transforming Mental Health Crisis Care
Published by the State of Queensland (Queensland Health), December 2020

- There is a need to develop a data-driven quality improvement approach to inform clinical, cultural and system changes that will lead to improved outcomes for people in crisis on the Gold Coast.
- There is a need to add to the local evidence base through research and evaluation of crisis reform initiatives.
- New models of service to respond to mental health crisis will require training and support to ensure success.
- With lived experience workers a central component of the crisis service models, there are specific needs to ensure enough peer workers and appropriate support systems are in place.
- Opportunities for shared training across organizations can assist in achieving a consistent approach and shared language, attitudes and beliefs which will be important in an integrated network of care and include:
 - training developed to support all underlying principles and new models of service
 - opportunities to enhance connections of networks of services through shared training
 - lived experience workforce training
 - all staff receive training on crisis intervention, trauma-informed care, lived experience principles.

Service system

Services	Number in GCPHN region	Distribution	Capacity discussion
GCPHN funded Psychological Services Program (PSP) suicide prevention	Of the 20 contracted organisations, 16 are contracted to provide suicide prevention services	Providers are distributed across the region	Dedicated suicide prevention services on the Gold Coast appear to be limited; however, some mental health services provide information and referral advice on suicide prevention.
Gold Coast Health crisis helpline	1 phone hotline (13 MH CALL) for the Acute Care Treatment (ACT) Team	ACT team telephone service available 24hrs	
Emergency Departments (ED)	5	Southport and Robina (public) Southport, Benowa and Tugun (private)	A 2018 review of clients accessing Psychological Services Program (PSP) suicide prevention service stream indicates strong use but those using the service tend to be females and younger people, which are not the most at risk cohorts in the region.
Support and Transition Program - Suicide Prevention (coordination support for those at-risk of suicide, recently attempted or are recently discharged)	1	Accessible via contact with public hospitals in Robina and Southport	
Crisis helplines	4 national (Lifeline, Suicide Call Back Service, Mensline, Kids Helpline)	24-hour, 7-day telephone services. Public knowledge of these services would drive uptake/demand.	Crisis services on the Gold Coast are available through the public health system in the form of hospital emergency departments and specific crisis support (Acute Care Treatment team, 24hr phone line).
Counselling helplines and websites	12 national help lines (Mensline, Kids Helpline, Open Arms formerly Veterans and Veterans Families Counselling Service, QLife, Carers Australia, eheadspace, 1800 Respect, Relationships Australia, SANE Australia, ReachOut, BeyondBlue, Counselling Online, Child abuse prevention service)	Online and telephone services.	There are numerous well-known national suicide prevention and crisis services that are likely to be accessed by the Gold Coast community. For example, Lifeline (phone and online), Suicide call back service (phone and online) and Beyond blue (phone and online).

			There are no specialised suicide prevention or crisis services for Aboriginal and Torres Strait Islander people on the Gold Coast although the Acute Care Team does employ an Aboriginal and Torres Strait Islander Mental Health Worker.
--	--	--	---

Consultation

GCPHN and Gold Coast Health have been working collaboratively over the past two years through the development of three separate but complementary regional strategies: Joint Regional Plan for Mental Health, Suicide Prevention, Alcohol and Other Drugs Services (JRP), Crisis Reform Strategy (CRS), and the Suicide Prevention Community Action Plan (SP CAP). While there are different drivers for each of these strategies, many of the underlying issues and longer-term outcomes are similar resulting in the interrelated nature of the three strategies and their contributing activities.

A range of consultation activities have occurred over the past two years to support the development of these strategies as well as the needs assessment process, service design, implementation and evaluation. Community members, clinicians and service providers have been engaged through various mechanisms including workshops, advisory groups, consumer journey mapping, one-to-one interviews, sector presentations, working groups and co-design processes. In addition to the findings mentioned above aligned to the LifeSpan framework, the following insights have emerged from consultation activities:

Service provider consultation

- People presenting to hospital feeling at risk of self-harm but whose mental health issues are not seen as serious enough for admission with limited follow up provided.
- Training and skills development for school staff will support early identification, intervention and referrals.
- Need for enhancing the skills of mainstream services, GPs, and clinicians to work with at risk and vulnerable populations.
- Limited community support systems and services available for those that have attempted suicide
- Early identification of at-risk people who identify as LGBTIQAP+ is key to suicide prevention.
- Lotus staff have described emerging impacts of COVID-19 on service delivery. This includes increased number of people requiring supports and connections to Centrelink and additional time required to support clients in the use of technology to facilitate connections and access to other services and supports during this time.
- The Social and Economic fallout of COVID-19 is anticipated to have significant impacts on service demand and need.

- Responses for 45-56-year age demographic remains a definite gap. People are left highly vulnerable due to unplanned/unforeseen circumstances with little support from the community.
- Access to Domestic Violence services have been an issue especially with carers and violence orders, gaps evident and challenges with this sector.
- GPs refer to the PSP program on “need”, usually distress rather than personal attributes (such as being LGBTIQAP+ or CALD).
- The GCPHN region would benefit greatly from a Safe Space/Safe Haven Service located within walking distance to one of the major hospitals such as Robina.
- The current After Hours Model is limited in its reach (location wise) and capacity to focus around crisis intervention/ED presentation reduction rather than a broader catchment model which includes much of homelessness supports/drop ins.

Service user consultation

- Inadequate response for individuals presenting to hospital feeling unsafe/at risk of self-harm but who are not admitted as their immediate health issues are not seen as serious or acute enough.
- Limited community support systems or services for those that have attempted suicide.
- People who have survived suicide attempts want more support, particularly with non-health related issues such as financial support, relationships and housing.
- Individuals being discharged feel excluded from the hospital discharge planning process.
- Due to high numbers of persons presenting with high mental health needs and/or risk of suicide there are periods of increased length of response times from the Acute Care Team.
- When describing their experience of care, consumers frequently express a lack of empathy and compassion from primary care providers.



Australian Government



An Australian Government Initiative

“Building one world class health system for the Gold Coast.”

Gold Coast Primary Health Network
Primary Care Gold Coast Limited (ABN 47 152 953 092), trading as Gold Coast Primary Health Network.
All rights reserved. Content is correct at time of publication.

Level 1, 14 Edgewater Court Robina QLD 4226 | PO Box 3576, Robina Town Centre QLD 4230
P 07 5635 2455 | E info@gcphn.com.au | W www.gcphn.org.au

Gold Coast Primary Health Network (GCPHN) gratefully acknowledges the financial and other support from the Australian Government Department of Health.



Gold Coast Primary Health Network would like to acknowledge and pay respect to the land and the traditional practices of the families of the Yugambeh Language Region of South East Queensland and their Elders past, present and emerging.

