

# Clinical Age limits for vaccine usage (not fundingrelated)

**ATAGI** (Australian Technical Advisory Group on Immunisation) - Recommended

**PI** (product information)

*\*Please refer to Variations from Product information found in each specific vaccine preventable disease.*

<b>Act-HIB and Hiberix</b>	For use from $\geq 6$ weeks to $< 5$ years of age: PI For use from $\geq 6$ weeks to $< 5$ years of age and older people with asplenia people who have received an allogeneic or autologous hematopoietic stem cell transplant (ATAGI)
<b>Adacel</b>	For use in $\geq 10$ years of age.
<b>ADT</b>	For use in $\geq 10$ years of age (primary or booster): ATAGI
<b>Bexsero</b>	From 6 weeks of age: ATAGI
<b>Boostrix</b>	For use in people $\geq 10$ years of age, except in certain circumstances. ATAGI
<b>Engerix -B (paediatric)</b> <b>H-B-Vax (paediatric)</b>	For use from birth to $< 20$ years of age. See (adult vaccine) for $11 \leq 15$ years of age.
<b>Engerix B (adult)</b> <b>H-B-VaxII (adult)</b>	For use from $11 \leq 15$ years of age. For use from $\geq 20$ years of age.
<b>Gardasil</b>	Recommended for males aged $> 26$ years in high-risk groups, such as men who have sex with men, and people of any age who are immunocompromised. Females aged 9 to $< 46$ years and males aged 9 to $< 27$ years: PI
<b>Havrix Junior</b>	For use in people aged $2 \leq 15$ years of age.
<b>Infanrix</b>	For use from 6 weeks to $< 10$ years of age.
<b>Infanrix -Hexa/ Vaxelis</b>	For use from 6 weeks to $< 10$ years of age.
<b>Infanrix-IPV</b>	For use from 6 week $< 10$ years of age.
<b>I POL</b>	For use from $\geq 6$ weeks of age.
<b>MMRII / Priorix</b>	For use from $\geq 12$ months of age. For use from $\geq 6$ months of age ATAGI (if travel to highly endemic areas) 2 further doses are required after 11 months of age: ATAGI
<b>Nimenrix</b>	For use from $\geq 6$ weeks.
<b>Prevenar13</b>	For use in children aged $\geq 6$ weeks and in adults.
<b>Pneumovax 23</b>	For use in children aged $\geq 2$ years and in adults.
<b>Priorix-Tetra</b>	For use $\geq 12$ months $< 14$ years of age.
<b>Quadracel</b>	For use from $\geq 6$ weeks of age to $< 10$ years: ATAGI
<b>Rotarix</b>	For use in infants $\geq 6$ weeks, dose 1 cut off 14 weeks and 6 days. Dose 2 cut off from 24 weeks and 6 days.
<b>Shingrix</b>	For use in over 18 immunocompromised adults. (refer to eligible groups) For use in Aboriginal and Torres Strait Islanders $\geq 50$ For use in adults $\geq 65$ years and older (non- Indigenous)
<b>Tripacel</b>	For use from $\geq 6$ weeks of age to $< 10$ years: ATAGI
<b>Vaqtia (paed)</b>	For use from $\geq 12$ months to $< 18$ years of age.
<b>Varilrix</b>	For use $\geq 9$ months: PI (not valid until $\geq 12$ months of age)
<b>Varivax</b>	For use in people aged $\geq 12$ months: PI (not valid until $\geq 12$ months of age)